## CLAIMS

 A pharmaceutical composition for the prevention or treatment of overactive bladder accompanied with neurogenic disorders, which comprises as an active ingredient an indoline derivative represented by a formula:

[Chem.1]

$$\begin{pmatrix}
H \\
N \\
R
\end{pmatrix}$$

$$\begin{pmatrix}
H \\
N \\
OH_3
\end{pmatrix}$$

$$OR^2$$
(1)

in the formula, R represents a saturated or unsaturated aliphatic 10 acyl group which may have as a substituent a hydroxy group, a lower alkoxy group, a carboxy group, a lower alkoxycarbonyl group, a cycloalkyl group, an aryl group or one or more halogen atoms; a hydroxy lower alkyl group; an aliphatic acyloxyalkyl group; a lower alkyl group which has as a substituent a lower alkoxy group, a carboxy group, a lower alkoxycarbonyl group, an 15 aryl-substituted lower alkoxycarbonyl group, a carbamoyl group, a mono or di (lower alkyl)-substituted carbamoyl group or a cyano group; an aromatic acyl group which may have as a substituent one or more halogen atoms; a furoyl group or a pyridylcarbonyl group; R1 represents a cyano group or a carbamoyl group; R2 20 represents a lower alkyl group which may have as a substituent a cyano group, an aryl group or one or more halogen atoms; or a pharmaceutically acceptable salt thereof.

25 2. A pharmaceutical composition as claimed in claim 1 wherein the active ingredient is (-)-1-(3-hydroxypropy1)-5-

- ((2R)-2-{[2-({2-[(2,2,2-trifluoroethyl)oxy]phenyl}oxy)-ethyl]amino}propyl)-2,3-dihydro-1H-indol-7- carboxamide or a pharmaceutically acceptable salt thereof.
- 5 3. A pharmaceutical composition as claimed in claim 1 or 2 wherein the neurogenic disorder is cerebral infarction, Parkinson's disease, spinal cord involvement, peripheral neurogenic disorder or multiple sclerosis.
- 10 4. A pharmaceutical composition as claimed in any of claims 1 to 3, which is used in combination with one or more other agents used for overactive bladder accompanied with neurogenic disorder.
- 5. A pharmaceutical composition as claimed in claim 4 wherein the other agent used for overactive bladder accompanied with neurogenic disorder is an agent selected from an anticholinergic drug, an anti-anxiety drug, a cholinergic drug, a cholinestersse inhibitor, an antispasmodic drug, an anti-inflammatory drug and an antimicrobial drug.